

32-7614: Recombinant Mouse Cell Adhesion Molecule 1/CADM1/IGSF4A/SynCAM1 (C-Fc)

Gene : Cadm1
Gene ID : 54725
Uniprot ID : Q8R5M8

Description

Source: Human Cells.
MW :65.3kD.

Recombinant Mouse Cell adhesion molecule 1 is produced by our Mammalian expression system and the target gene encoding Gln48-His388 is expressed with a Fc tag at the C-terminus. Cell adhesion molecule 1(CADM1) is a single-pass type I membrane protein and belongs to the nectin family. It contains 2 Ig-like C2-type (immunoglobulin-like) domains and 1 Ig-like V-type (immunoglobulin-like) domain. CADM1 acts as a tumor suppressor in non-small-cell lung cancer (NSCLC) cells. Interaction with CRTAM promotes natural killer (NK) cell cytotoxicity and interferon-gamma (IFN-gamma) secretion by CD8+ cells in vitro as well as NK cell-mediated rejection of tumors expressing CADM3 in vivo. CADM1 may contribute to the less invasive phenotypes of lepidic growth tumor cells. In mast cells, it may mediate attachment to and promote communication with nerves. CADM1, together with MITF, is essential for development and survival of mast cells in vivo. The protein acts as a synaptic cell adhesion molecule and plays a role in the formation of dendritic spines and in synapse assembly. It may be involved in neuronal migration, axon growth, pathfinding, and fasciculation on the axons of differentiating neurons. CADM1 may play diverse roles in the spermatogenesis including in the adhesion of spermatocytes and spermatids to Sertoli cells and for their normal differentiation into mature spermatozoa.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : QNLFTKDVTVIEGEVATISCVNKSDDSVIQLLNPNRQTIYFRDFRPLKDSRFQLLNFSSELKVSLTNVSIISDEGRYFCQLYTDPPQESYTTITVLVPPRNLMIQKDTAVEGEEIEVNCTAMASKPATTIRWFKGNKELKGKSEVEEWSDMYTVTSQMLKVKHEDDGVVICQVEHPAVTGNLQTRYLEVQYKPVHMQMTYPLQGLTREGDAFELTCEAIGKPQPMVMTWVRVDDDEMPQHAVLSGPNLFINNLNKTNDNGTYRCEASNIVGKAHSDYMLYVYDPPPTTIPPTTTTTTTTTTTTTILITDITATTEPAVHDSRAGEEGTIGAVDHVDDIEGRMDEPKSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.