

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7617: Recombinant Human Teratocarcinoma-Derived Growth Factor 1/TDGF1/Cripto (C-Fc)(Discontinued)

Gene ID: TDGF1
Gene ID: 6997
Uniprot ID: P13385

## **Description**

Source: Human Cells.

MW:42.8kD.

Recombinant Human Teratocarcinoma-derived growth factor 1 is produced by our Mammalian expression system and the target gene encoding Leu31-Ser169 is expressed with a Fc tag at the C-terminus. Teratocarcinoma-derived growth factor 1(TDGF1) is a Cell membrane protein and contains 1 EGF-like domain. The protein plays an essential role in embryonic development and tumor growth. Mutations in this gene are associated with forebrain defects. It also may play a role in the determination of the epiblastic cells that subsequently give rise to the mesoderm.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

**Content:** Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

Lyophilized protein should be stored at -20  $^{\circ}$ C, though stable at room temperature for 3 weeks.

**Storage condition:** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: LGHQEFARPSRGYLAFRDDSIWPQEEPAIRPRSSQRVPPMGIQHSKELNRTCCLNGGTCMLGSFCACPPSFYG

RNCEHDVRKENCGSVPHDTWLPKKCSLCKCWHGQLRCFPQAFLPGCDGLVMDEHLVASRTPELPPSVDDIEG RMDEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVH NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMT KNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL

HNHYTQKSLSLSPGK

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin**: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.