

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7628: Recombinant Mouse Dickkopf-Related Protein 1/mDKK1/DKK-1 (N-8His)(Discontinued)

**Gene ID:** Dkk1 **Gene ID:** 13380 **Uniprot ID:** 054908

## **Description**

Source: Human Cells.

MW:27.4kD.

Recombinant Mouse Dickkopf-related protein 1 is produced by our Mammalian expression system and the target gene encoding Thr32-His272 is expressed with a 8His tag at the N-terminus. This gene encodes a protein that is a member of the dickkopf family. It possesses two clusters of ten cysteine residues separated by a linker region. The protein highly expressed in thyroid, small intestine, stomach, liver, placenta, pancreas, uterus, abdominal cavity, bladder and skin. Weaker expression has been detected in colon and spleen. It antagonizes canonical Wnt signaling by inhibiting LRP5/6interaction with Wnt and by forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6.

## **Product Info**

**Amount:** 10 μg / 50 μg

**Content:** Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

Lyophilized protein should be stored at -20  $^{\circ}$ C, though stable at room temperature for 3 weeks.

**Storage condition:** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: HHHHHHHHHQTLNSVLINSNAIKNLPPPLGGAGGQPGSAVSVAPGVLYEGGNKYQTLDNYQPYPCAEDEECGS

DEYCSSPSRGAAGVGGVQICLACRKRRKRCMRHAMCCPGNYCKNGICMPSDHSHFPRGEIEESIIENLGNDHN AAAGDGYPRRTTLTSKIYHTKGQEGSVCLRSSDCAAGLCCARHFWSKICKPVLKEGQVCTKHKRKGSHGLEIF

QRCYCGEGLACRIQKDHHQASNSSRLHTCQRH

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than  $0.1 \text{ ng}/\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.