

32-7683: Recombinant Human Fumarylacetoacetase/FAH (C-6His)

Gene : FAH
Gene ID : 2184
Uniprot ID : P16930

Description

Source: Human Cells.
MW :47.4kD.

Recombinant Human Fumarylacetoacetase is produced by our Mammalian expression system and the target gene encoding Ser2-Ser419 is expressed with a 6His tag at the C-terminus. Fumarylacetoacetase belongs to the FAH family. Fumarylacetoacetase is primarily expressed in liver and kidney. It exists as a homodimer and catalyzes the hydrolysis of 4-fumarylacetoacetate into fumarate and acetoacetate. Defects in Fumarylacetoacetase cause tyrosinemia type 1, which is a congenital metabolism defect characterized by elevated levels of tyrosine in the blood and urine, and hepatorenal manifestations. Typical features include renal tubular injury, self-mutilation, hepatic necrosis, episodic weakness, and seizures.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : SFIPVAEDSDFPIHNLPGYVGFSTRGDPRPRIGVAIGDQILDLSIIKHLFTGPVLSKHQDVFNQPTLNSFMGLGQAA WKEARVFLQNLLSVSQARLRDDTELKCAFISQASATMHLPATIGDYTDYSSRQHATNVGIMFRDKENALMP NWLHLPVGYHGRASSVVVSGTPIRRPMGQMKPDDSKPPYVYACKLLDMELEMAFFVGPNGRLGEPISKAE HIFGMVLMNDWSARDIQKWEYVPLGPFGLGKSFVTVSPWVVPMDALMPFAVNPQDPRPLPYLCHDEPYTF DINLSVNLKGEQMSQAATICKSNFKYMYWTMLQQLTHHSVNGCNLRPGDLLASGTISGPEPEPFGSMLELSWK GTKPIDLGNQTRKFLLDGDEVIITGYCQGDGYRIGFGQCAGKVLPAALLPSVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.