

32-7686: Recombinant Human Alcohol Dehydrogenase Class 4 Mu/ADH7 (C-6His)

Gene : ADH7
Gene ID : 131
Uniprot ID : P40394

Description

Source: Human Cells.
MW :42.5kD.

Recombinant Human ADH7 is produced by our Mammalian expression system and the target gene encoding Met1-Phe386 is expressed with a 6His tag at the C-terminus. Alcohol dehydrogenase class 4 mu/sigma chain (ADH7) is a cytoplasm enzyme which is a member of the alcohol dehydrogenase family. The expression of this gene makes it much more abundant in the stomach than the liver, thus it differs from the other known gene family members. ADH7 may participate in the synthesis of retinoic acid, a hormone important for cellular differentiation. Medium-chain (octanol) and aromatic (m-nitrobenzaldehyde) compounds are the best substrates. Ethanol is not a good substrate but at the high ethanol concentrations reached in the digestive tract, it plays a role in the ethanol oxidation and contributes to the first pass ethanol metabolism.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS,PH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : MFAEIQDKDRMGTAGKVIKCKAAVLWEQKQPFSEIEVAPPKTKEVRIKILATGICRTDDHVIKGMTVSKFPVI
VGHEATGIVESIGEGVTTVKPGDKVIPLFPQCRECNACRNPDGNLCSITGRGVLADGTTFRFTCKGKPVVHHF
MNTSTFTFTEYTVVDESSVAKIDDAAPPEKVKCLIGCGFSTGYGAAVKTKGKVKPGSTCVVFGLGGVGLSVIMGCKSA
GASRIIGIDLNKDKFEKAMAVGATECISPKDSTKPISEVLSEMTGNNVGYTFEVIHLETMIDALASCHMNYGTSV
VVGVPSSAKMLTYDPMMLLFTGRTWKGCVFGGLKSRDDVPKLVTEFLAKKFDLQDLITHVLPFKKISEGFELLNSG
QSIRTVLTFVDHHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.