

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7697: Recombinant Human Netrin-G1/NTNG1 (C-6His)

Gene ID: 22854 Uniprot ID: Q9Y212

Description

Source: Human Cells.

MW:43.5kD.

Recombinant Human Netrin-G1 is produced by our Mammalian expression system and the target gene encoding His29-Ser409 is expressed with a 6His tag at the C-terminus. Netrin-G1 (NTNG1) is a member of a conserved family of proteins that act as axon guidance cues during vertebrate nervous system development. Netrin-G1 contains one laminin EGF-like domain and one laminin N-terminal domain, Netrin-G1 is highly expressed in the thalamus, lowly in other tissue. Netrin-G1 localizes to the cell membrane. Netrin-G1 interacts with NGL1 and is glycosylated in the N-terminal. In addition, Netrin-G1 can promotesneurite outgrowth of both axons and dendrites.

Product Info

Amount: 10 μg / 50 μg

Content: Lyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: HYDLCKTQIYTEEGKVWDYMACQPESTDMTKYLKVKLDPPDITCGDPPETFCAMGNPYMCNNECDASTPELAH

PPELMFDFEGRHPSTFWQSATWKEYPKPLQVNITLSWSKTIELTDNIVITFESGRPDQMILEKSLDYGRTWQPYQ YYATDCLDAFHMDPKSVKDLSQHTVLEIICTEEYSTGYTTNSKIIHFEIKDRFAFFAGPRLRNMASLYGQLDTTKK LRDFFTVTDLRIRLLRPAVGEIFVDELHLARYFYAISDIKVRGRCKCNLHATVCVYDNSKLTCECEHNTTGPDCG KCKKNYQGRPWSPGSYLPIPKGTANTCIPSISSIGTNVCDNELLHCQNGGTCHNNVRCLCPAAYTGILCEKLRCE

EAGSCGSVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.