

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7702: Recombinant Human CEACAM7/CGM2 (C-6His)

Gene ID: CEACAM7
Gene ID: 1087
Uniprot ID: 014002

Description

Source: Human Cells.

MW:13.1kD.

Recombinant Human CEACAM7 is produced by our Mammalian expression system and the target gene encoding Thr36-Phe142 is expressed with a 6His tag at the C-terminus. Carcinoembryonic Antigen-Related Cell Adhesion Molecule 7 (CEACAM7) is a member of the immunoglobulin superfamily A and CEA family. CEACAM7 localizes to the cell membrane and contains one Ig-like C2-type domain and one Ig-like V-type domain. The expression of CEACAM7 is significantly decreased in rectal cancer. Differences in CEACAM7 expression levels between long-term survivors and those with recurrent disease introduce a potential tumor marker to define a subset of patients who benefit most from adjuvant therapy.

Product Info

Amount: 10 μg / 50 μg

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: TNIDVVPFNVAEGKEVLLVVHNESQNLYGYNWYKGERVHANYRIIGYVKNISQENAPGPAHNGRETIYPNGTLLI

QNVTHNDAGIYTLHVIKENLVNEEVTRQFYVFVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.