

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7704: Recombinant Human Serpin A4/Kallistatin (C-6His)

Gene ID: SERPINA4
Gene ID: 5267
Uniprot ID: P29622

## **Description**

Source: Human Cells.

MW:47.1kD.

Recombinant Human Serpin A4 is produced by our Mammalian expression system and the target gene encoding Gln21-Pro427 is expressed with a 6His tag at the C-terminus. Serpin Peptidase Inhibitor, Clade A (a-1 Antiproteinase, Antitrypsin), Member 4 (Serpin A4) is a member of the Serpin family. Serpin A4 exists as a monomer and some homodimers. Serpin A4 is expressed by the liver and secreted in plasma. Serpin A4 is a regulator of vascular homeostasis capable of controlling a wide spectrum of biological actions in the cardiovascular and renal systems. It can inhibit intracellular reactive oxygen species formation in cultured cardiac and renal cells. In addition, Serpin A4 has anti-inflammatory effect. Heparin blocks kallistatin's complex formation with tissue kallikrein and abolishes its inhibitory effect on tissue kallikrein's activity.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of 50mM TrisHCl, 150mM NaCl, 10mM NaCl, pH 8.0.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: QLHVEHDGESCSNSSHQQILETGEGSPSLKIAPANADFAFRFYYLIASETPGKNIFFSPLSISAAYAMLSLGACSH

SRSQILEGLGFNLTELSESDVHRGFQHLLHTLNLPGHGLETRVGSALFLSHNLKFLAKFLNDTMAVYEAKLFHTN FYDTVGTIQLINDHVKKETRGKIVDLVSELKKDVLMVLVNYIYFKALWEKPFISSRTTPKDFYVDENTTVRVPMML QDQEHHWYLHDRYLPCSVLRMDYKGDATVFFILPNQGKMREIEEVLTPEMLMRWNNLLRKRNFYKKLELHLPK FSISGSYVLDQILPRLGFTDLFSKWADLSGITKQOKLEASKSFHKATLDVDEAGTEAAAATSFAIKFFSAQTNRHI

LRFNRPFLVVIFSTSTQSVLFLGKVVDPTKPHHHHHH

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than  $0.1 \text{ ng}/\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.