

32-7715: Recombinant Human Isocitrate Dehydrogenase 1/IDH1 (C-6His)

 Gene :
 IDH1

 Gene ID :
 3417

 Uniprot ID :
 075874

Description

Source: Human Cells.

MW :47.7kD.

Recombinant Human Isocitrate Dehydrogenase is produced by our Mammalian expression system and the target gene encoding Met1-Leu414 is expressed with a 6His tag at the C-terminus. Isocitrate Dehydrogenase [NADP] Cytoplasmic (IDH1) belongs to the isocitrate and isopropylmalate dehydrogenases family. IDH1 exists as a homodimer, binding one magnesium or manganese ion per subunit. Mutations of IDH1 have been shown to cause metaphyseal chondromatosis with aciduria and are involved in the development of glioma IDH plays a role in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2oxoglutarate, namely the a-hydroxylation of phytanic acid.

Product Info

Amount :	10 μg / 50 μg
Content :	Supplied as a 0.2 μ m filtered solution of PBS.
Storage condition :	Store at -20°C, stable for 6 months after receipt. Please minimize freeze-thaw cycles.
Amino Acid :	MSKKISGGSVVEMQGDEMTRIIWELIKEKLIFPYVELDLHSYDLGIENRDATNDQVTKDAAEAIKKHNVGVKCAT ITPDEKRVEEFKLKQMWKSPNGTIRNILGGTVFREAIICKNIPRLVSGWVKPIIIGRHAYGDQYRATDFVVPGPGK VEITYTPSDGTQKVTYLVHNFEEGGGVAMGMYNQDKSIEDFAHSSFQMALSKGWPLYLSTKNTILKKYDGRFK DIFQEIYDKQYKSQFEAQKIWYEHRLIDDMVAQAMKSEGGFIWACKNYDGDVQSDSVAQGYGSLGMMTSVLV CPDGKTVEAEAAHGTVTRHYRMYQKGQETSTNPIASIFAWTRGLAHRAKLDNNKELAFFANALEEVSIETIEAGF MTKDLAACIKGLPNVQRSDYLNTFEFMDKLGENLKIKLAQAKLVDHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/Ã^{\[}µg (1 IEU/Ã[[]µg) as determined by LAL test.