

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7718: Recombinant Human Immunoglobulin Superfamily Member 8/IGSF8/CD316 (C-6His)

**Gene ID:** 93185 **Uniprot ID:** 9969P0

## **Description**

Source: Human Cells.

MW:59.6kD.

Recombinant Human IGSF8 is produced by our Mammalian expression system and the target gene encoding Arg28-Thr579 is expressed with a 6His tag at the C-terminus. Immunoglobulin Superfamily Member 8 (IGSF8) is a single-pass membrane protein. IGSF8 contains four Ig-like C2 type domains. The Ig-like C2-type domains 3 and 4 are required for interactions with CD81. IGSF8 may regulate proliferation and differentiation of keratinocytes. IGSF8 may participate in the regulation of neurite outgrowth and maintenance of the neural network in the adult brain. It also may play a role on integrin-dependent morphology and motility functions.

## **Product Info**

**Amount:** 10 μg / 50 μg

**Content:** Lyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: REVLVPEGPLYRVAGTAVSISCNVTGYEGPAQQNFEWFLYRPEAPDTALGIVSTKDTQFSYAVFKSRVVAGEVQ

VQRLQGDAVVLKIARLQAQDAGIYECHTPSTDTRYLGSYSGKVELRVLPDVLQVSAAPPGPRGRQAPTSPPRM TVHEGQELALGCLARTSTQKHTHLAVSFGRSVPEAPVGRSTLQEVVGIRSDLAVEAGAPYAERLAAGELRLGKE GTDRYRMVVGGAQAGDAGTYHCTAAEWIQDPDGSWAQIAEKRAVLAHVDVQTLSSQLAVTVGPGERRIGPG EPLELLCNVSGALPPAGRHAAYSVGWEMAPAGAPGPGRLVAQLDTEGVGSLGPGYEGRHIAMEKVASRTYRL RLEAARPGDAGTYRCLAKAYVRGSGTRLREAASARSRPLPVHVREEGVVLEAVAWLAGGTVYRGETASLLCNI SVRGGPPGLRLAASWWVERPEDGELSSVPAQLVGGVGQDGVAELGVRPGGGPVSVELVGPRSHRLRLHSLG

PEDEGVYHCAPSAWVQHADYSWYQAGSARSGPVTVYPYMHALDTVDHHHHHH

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \Box \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin**: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.