

32-7718: Recombinant Human Immunoglobulin Superfamily Member 8/IGSF8/CD316 (C-6His)

Gene : IGSF8
Gene ID : 93185
Uniprot ID : Q969P0

Description

Source: Human Cells.
MW :59.6kD.

Recombinant Human IGSF8 is produced by our Mammalian expression system and the target gene encoding Arg28-Thr579 is expressed with a 6His tag at the C-terminus. Immunoglobulin Superfamily Member 8 (IGSF8) is a single-pass membrane protein. IGSF8 contains four Ig-like C2 type domains. The Ig-like C2-type domains 3 and 4 are required for interactions with CD81. IGSF8 may regulate proliferation and differentiation of keratinocytes. IGSF8 may participate in the regulation of neurite outgrowth and maintenance of the neural network in the adult brain. It also may play a role on integrin-dependent morphology and motility functions.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.2.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : REVLVPEGPLYRVAGTAVSISCNVTGYEGPAQQNFEWFLYRPEAPDTALGIVSTKDTQFSYAVFKSRVVAGEVQ VQRLQGDAVVLKIARLQAQDAGIYECHTPTDTRYLGYSYSGKVELRVLPDVLQVSAAPPGPRGRQAPTSPRM TVHEGQELALGCLARTSTQKHTHLAVSFGRSVPEAPVGRSTLQEVVGIRSDLAVEAGAPYAERLAAGELRLGKE GTDRYRMVVGGAQAGDAGTYHCTAAEWIQDPDGSWAQIAEKRAVLAHVVDVQTLSSQLAVTVGPGERRIGPG EPLELLCNVSGALPPAGRHAAYSVGWEMAPAGAPGPGRLVAQLDTEGVGSLGPGYEGRHIAMEKVASRTYRL RLEAARPGDAGTYRCLAKAYVRGSGTRLREAASARSRPLPVHVREEGVVLEAVAWLAGGTVYRGETASLLCNI SVRGGPPGLRLAASWWWVERPEDGELSSVPAQLVGGVQDQVAELGVRPGGGPPSVELVGP RSHRLRLHSLG PEDEGVYHCAPSAWVQHADYSWYQAGSARSGPVTYYPYMHALDVTVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.