

32-7728: Recombinant Human Leucine-Rich Repeat Transmembrane Protein FLRT3/FLRT3 (C-6His)

Gene : FLRT3
Gene ID : 23767
Uniprot ID : Q9NZU0

Description

Source: Human Cells.

MW :57.8kD.

Recombinant Human FLRT3 is produced by our Mammalian expression system and the target gene encoding Lys29-Pro528 is expressed with a 6His tag at the C-terminus. Leucine-Rich Repeat Transmembrane Protein FLRT3 (FLRT3) is a member of the fibronectin leucine rich transmembrane protein (FLRT) family. Proteins in this family play an role in cell adhesion and/or receptor signalling. FLRT3 is a single-pass type I membrane protein and contains one fibronectin type-III domain, ten LRR (leucine-rich) repeats, one LRRCT domain, and one LRRNT domain. FLRT3 may have a function in cell adhesion and/or receptor signaling. FLRT3 may regulate cellular adhesion between the epithelial apical ridge and the underlying mesenchyme and in establishing the dorso-ventral position of the ridge.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.2.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : KSCPSVCRCDAGFIYCNDRFLTSIPTGIPEDATTLYLQNNQINNAGIPSDLKKNLLKVERIYLYHNSLDEFPTNLPKY VKELHLQENNIRITITYDSLKIPYLEELHLDNSVSAVSIEEGAFRDSNYLRLFLSRNHLSTIPWGLPRTIEELRLD DNRISTISSPSLQGLTSLKRLVLDGNLLNNHGLGDKVFFNLVNLTELSLVRNSLTAAPVNLPGTNLRKLYLQDNHI NRVPPNAFSYLRQLYRLDMSNNNLSNLPQGIFDDLDNITQLILRNNPWYCGCKMKWVRDWLQSLPVKVNVRG LMCQAPEKVRGMAIKDLNAELFDCKDSGIVSTIQITTAIPNTVYPAQGQWPAPVTKQPDIKNPKLTKDHQTTGSP SRKTITITVKSVTSDTIHISWKLALPMTALRLSWLKLGHSPAFGSITETIVTGERSEYLVTALEPDSPYKVCMPME TSNLYLFDETPVCIEETETAPLRMYNPTTTLNREQEKEPYKNPNLPVDHHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.