

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7731: Recombinant Human SH3KBP1/CD2BP3/CIN85v(C-6His)(Discontinued)

Gene: SH3KBP1 Gene ID: 30011 Uniprot ID: Q96B97

Description

Source: Human Cells.

MW:74.1kD.

Recombinant Human SH3KBP1 is produced by our Mammalian expression system and the target gene encoding Met1-Lys665 is expressed with a 6His tag at the C-terminus. SH3 Domain-Containing Kinase-Binding Protein 1 (SH3KBP1) contains 3 SH3 domains. SH3KBP1 is widely expressed in many tissues, it is also expressed in some cancer cell lines. SH3KBP1 plays a role in the regulation of endocytosis and lysosomal degradation of ligand-induced receptor tyrosine kinases. SH3KBP1 is involved in regulation of ligand-dependent endocytosis of the IgE receptor and in the regulation of cellular stress response via the MAPK pathways. SH3KBP1 is also required for the control of cell shape and migration.

Product Info

Amount: $10 \mu g / 50 \mu g$

Lyophilized from a 0.2 µm filtered solution of 50mM Tris-HCI,10mM reduced Content:

Glutathione,pH=8.0.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Storage condition:

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: MVEAIVEFDYQAQHDDELTISVGEIITNIRKEDGGWWEGQINGRRGLFPDNFVREIKKEMKKDPLTNKAPEKPL

> HEVPSGNSLLSSETILRTNKRGERRRRRCQVAFSYLPQNDDELELKVGDIIEVVGEVEEGWWEGVLNGKTGMF PSNFIKELSGESDELGISQDEQLSKSSLRETTGSESDGGDSSSTKSEGANGTVATAAIQPKKVKGVGFGDIFKDK PIKLRPRSIEVENDFLPVEKTIGKKLPATTATPDSSKTEMDSRTKSKDYCKVIFPYEAQNDDELTIKEGDIVTLINKD CIDVGWWEGELNGRRGVFPDNFVKLLPPDFEKEGNRPKKPPPPSAPVIKQGAGTTERKHEIKKIPPERPEMLPN RTEEKERPEREPKLDLQKPSVPAIPPKKPRPPKTNSLSRPGALPPRRPERPVGPLTHTRGDSPKIDLAGSSLSGILDKDLSDRSNDIDLEGFDSVVSSTEKLSHPTTSRPKATGRRPPSQSLTSSSLSSPDIFDSPSPEEDKEEHISLAHRG VDASKKTSKTVTISQVSDNKASLPPKPGTMAAGGGGPAPLSSAVPSPLSSSLGTAGHRANSPSLFGTEGKPKME PAASSQAAVEELRTQVRELRSIIETMKDQQKREIKQLLSELDEEKKIRLRLQMEVNDIKKALQSKVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 $ng/\mu g$ (1 $IEU/\mu g$) as determined by LAL test.