

32-7768: Recombinant Human Linker for Activation of T-Cells Family Member 2/LAT2 (C-6His)

Gene : LAT2
Gene ID : 7462
Uniprot ID : Q9GZY6

Description

Source: Human Cells.
MW :25kD.

Recombinant Human Non-T cell activation linker is produced by our Mammalian expression system and the target gene encoding Arg27-Ala243 is expressed with a 6His tag at the C-terminus. Linker for Activation of T-Cells Family Member 2 (LAT2) is a single-pass type III membrane protein. LAT2 is highly expressed in the spleen, peripheral blood lymphocytes, and germinal centers of lymph nodes. LAT2 is involved in FCER1 (high affinity immunoglobulin epsilon receptor)-mediated signaling in mast cells. It may also be involved in BCR (B-cell antigen receptor)-mediated signaling in B-cells and FCGR1 (high affinity immunoglobulin gamma Fc receptor I)-mediated signaling in myeloid cells. Coupleing activate of these receptors and their associated kinases with distal intracellular events through the recruitment of GRB2.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : RCSRPGAKRSEKIYQQRSLREDQQSFTGSRTYSLVGQAWPGPLADMAPTRKDKLLQFYPSLEDPASSRYQNFSGSRHGSEEAYIDPIAMEYYNWGRFSKPPEDDDANSYENVLICKQKTTETGAQQEGIGGLCRGDLSSLALKTGPTSGLCPSASPEEDEESEDYQNSASIHQWRESRKVMGQLQREASPGVGPSPDEEDGEPDYVNGEVAATEAVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.