

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7785: Recombinant Human Activin Receptor 1B/Activin RIB/ALK-4/ACVR1B (C-6His)

Gene ID: ACVR1B
Gene ID: 91
Uniprot ID: P36896

Description

Source: Human Cells. MW:12.46kD.

Recombinant Human Activin Receptor IB is produced by our Mammalian expression system and the target gene encoding Ser24-Glu126 is expressed with a 6His tag at the C-terminus. Activin Receptor Type-1B (ACVR1B) is a single-pass type I membrane protein that belongs to the protein kinase superfamily. ACVR1B contains one GS domain and one protein kinase domain and is expressed in many tissues, most strongly in kidney, pancreas, brain, lung, and liver. ACVR1B acts as a transducer of activin or activin like ligands signals. Activin binds to either ACVR2A or ACVR2B and then forms a complex with ACVR1B, ACVR2A or ACVR2B activating ACVR1B through phosphorylation of its regulatory GS domain. They go on to recruit the R-SMADs, SMAD2 and SMAD3. ACVR1B also transducers signals of nodal, GDF-1, and Vg1. Mutations in ACVR1B are associated with pituitary tumors.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: SGPRGVQALLCACTSCLQANYTCETDGACMVSIFNLDGMEHHVRTCIPKVELVPAGKPFYCLSSEDLRNTHCC

YTDYCNRIDLRVPSGHLKEPEHPSMWGPVEVDHHHHHH

Application Note

Storage condition:

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.