

## 32-7786: Recombinant Human Natural Killer Cells Antigen CD94/CD94 (N-6His)

**Gene :** KLRD1  
**Gene ID :** 3824  
**Uniprot ID :** Q13241

### Description

Source: Human Cells.  
MW :17.7kD.

Recombinant Human Natural Killer Cells Antigen CD94 is produced by our Mammalian expression system and the target gene encoding Ser34-Ile179 is expressed with a 6His tag at the N-terminus. CD94 (Cluster of Differentiation 94), also known as killer cell lectin-like receptor subfamily D member 1 (KLRD1), is expressed on the surface of natural killer cells in the innate immune system. CD94 Plays a role as a receptor for the recognition of MHC class I HLA-E molecules by NK cells and some cytotoxic T-cells. CD94 Can form disulfide-bonded heterodimer with NKG2 family members. The CD94/NKG2 complex, on the surface of natural killer cells interacts with Human Leukocyte Antigen (HLA)-E on target cells. Natural killer (NK) cells are a distinct lineage of lymphocytes that mediate cytotoxic activity and secrete cytokines upon immune stimulation. Several genes of the C-type lectin superfamily, including members of the NKG2 family, are expressed by NK cells and may be involved in the regulation of NK cell function. KLRD1 (CD94) is an antigen preferentially expressed on NK cells and is classified as a type II membrane protein because it has an external C terminus.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** HHHHHHSFTKLSIEPAFTPGPNIELQKDSDCSCQEKWVGYRNCYFISSEQKTWNESRHLCASQKSSLLQLQ  
NTDELDFMSSSQFYWIGLSYSEEHTAWLWENGSAALSQYLFPSFETFNTKNCIAYNPNGNALDESCEDKNRYIC  
KQQLI

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.