

32-7794: Recombinant Human Transforming Growth Factor beta-1/TGFB1 (N-8His)

Gene : TGFB1
Gene ID : 7040
Uniprot ID : P01137

Description

Source: Human Cells.
MW :41.3&12.8kD.

Recombinant Human Transforming Growth Factor beta 1 is produced by our Mammalian expression system and the target gene encoding Leu30-Ser390 is expressed Transforming Growth Factor beta-1 (TGF beta-1) is a secreted protein which belongs to the TGF- beta family. TGF beta-1 is abundantly expressed in bone, articular cartilage and chondrocytes and is increased in osteoarthritis (OA). TGF beta-1 performs many cellular functions, including the control of cell growth, cell proliferation, cell differentiation and apoptosis. The precursor is cleaved into a latency-associated peptide (LAP) and a mature TGF beta-1 peptide. TGF beta-1 may also form heterodimers with other TGF beta family members. It has been found that TGF beta-1 is frequently upregulated in tumor cells. Mutations in this gene results in Camurati-Engelmann disease.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 50mM Glycine, 50mM NaCl, pH4.0.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : HHHHHHHHLLSTSKTIDMELVKKRIEAIKQILSKLRLASPPSQGEVPPGGLPEAVLALYNSTRDRVAGESAEPEPEPEADYYAKEVTRVLMVETHNEIYDKFKQSTHSIYMFNTSELREAVPEPVLLSRAELRLLRLKLVQHVLYQKYSNNSWRYLSNRLLAPSDSPEWLSFDVTGVVRQWLSRGGEIEGFRLSAHCSDSRDNTLQVDINGFTTGRRGDLATIHGMNRPFLLLMATPLERAQHLQSSRHRRALDTNYCFSSTEKNCCVRQLYIDFRKDLGWKWIHEPKGYPHANFCLGPCPYIWSLDTQYSKVLALYNQHNPGASAAPCCVPQALEPLPIVYVGRKPKVEQLSNMIVRSCKCS

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.