

## 32-7815: Recombinant Human NKG2-A/NKG2-B Type II Integral Membrane Protein(N-8His)

**Gene :** KLRC1  
**Gene ID :** 3821  
**Uniprot ID :** P26715

### Description

Source: Human cells.  
MW :16.5kD.

Recombinant Human NKG2-A/NKG2-B Type II Integral Membrane Protein is produced by our Mammalian expression system and the target gene encoding Arg100-Leu233 is expressed with a 8His tag at the N-terminus. NKG2-A/NKG2-B Type II Integral Membrane Protein contains 1 C-type lectin domain and belongs to the killer cell lectin-like receptor family. The killer cell lectin-like receptor family is a group of transmembrane proteins preferentially expressed in NK cells. Members of this proteins is characterized by the type II membrane orientation and the presence of a C-type lectin domain. NKG2 is expressed only in NK-cells, but not in T-cells or B-cells. It has been shown that NKG2 represents a family of related cDNA clones, designated NKG2A, NKG2B, NKG2C, and NKG2D, which encode type 2 integral membrane proteins (extracellular C-terminus) containing a C-type lectin domain. NKG2 plays a role as a receptor for the recognition of MHC class I HLA-E molecules by NK cells and some cytotoxic T-cells. NKG2A and NKG2B have been given the designation CD159a in the nomenclature of CD antigens.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** HHHHHHHHRHNNSSLNTRTQKARHCGHCPEEWITYSNSCYIGKERRTWEESLLACTSKNSSLLSIDNEEEMKFLSIIPSSWIGVFRNSSHHPWVTMNGLAFKHEIKSDNAELNCAVLQVNRLKSAQCGSSIIYHCKHKL

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.