

## 32-7822: Recombinant Human Parathyroid Hormone/PTH (N-8His)

**Gene :** PTH  
**Gene ID :** 5741  
**Uniprot ID :** P01270

### Description

Source: Human Cells.  
MW :10.5kD.

Recombinant Human Parathyroid Hormone is produced by our Mammalian expression system and the target gene encoding Ser32-Gln115 is expressed with a 8His tag at the N-terminus. Parathyroid hormone (PTH) is a critical hormone in the regulation of Ca<sup>++</sup> homeostasis. Parathyroid hormone is the most important endocrine regulator of calcium and phosphorus concentration in extracellular fluid. This hormone is secreted from cells of the parathyroid glands and finds its major target cells in bone and kidney. Another hormone, parathyroid hormone-related protein, binds to the same receptor as parathyroid hormone and has major effects on development. Like most other protein hormones, parathyroid hormone is synthesized as a prohormone. After intracellular processing, the mature hormone is packaged with in the Golgi into secretory vesicles, the secreted into blood by exocytosis. In renal epithelium, PTH promotes conversion of Vitamin D to its active form, lowers Ca<sup>++</sup> excretion and increases phosphate excretion. PTH also increases hematopoietic stem cell proliferation and mobilization and induces arterial vasodilation by regulating Ca<sup>++</sup> influx in PTH1R-expressing arterial smooth muscle.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** HHHHHHHHSVSEIQLMHNLGKHLNSMERVEWLRKKLQDVHNFVALGAPLAPRDAGSQRPKKEDNVLVESH  
EKSLGEADKADVNVLTAKASQ

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.