

32-7825: Recombinant Human Lymphocyte Function-associated Antigen 3/CD58/LFA-3 (C-Fc)(Discontinued)

Gene : CD58
Gene ID : 965
Uniprot ID : P19256

Description

Source: Human Cells.

MW :48.6kD.

Recombinant Human Lymphocyte function-associated antigen 3 is produced by our Mammalian expression system and the target gene encoding Phe29-Arg215 is expressed with a Fc tag at the C-terminus. Lymphocyte function-associated antigen 3 (LFA-3/CD58) is a single-pass type I membrane protein. CD58 is widely expressed on hematopoietic and non-hematopoietic human tissue and has been found on leukocytes, erythrocytes, endothelial cells, epithelial cells and fibroblasts of human origin. It is a Ligand of the T-lymphocyte CD2 glycoprotein. This interaction is important in mediating thymocyte interactions with thymic epithelial cells, antigen-independent and -dependent interactions of T-lymphocytes with target cells and antigen-presenting cells and the T-lymphocyte rosetting with erythrocytes. In addition, the LFA-3/CD2 interaction may prime response by both the CD2+ and LFA-3+ cells.

Product Info

Amount : 10 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4 .

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : FSQQIYGVVYGNVTFHVPSNVPLKEVLWKKQKDKVAELENSEFRAFSSFKNRVYLDTVSGSLTIYNLTSSDEDEY EMESPNITDTMKFFLYVLESPLSPTLTALTNLSIEVQCMIPPEYNSHRGLIMYSWDCPMEQCKRNSTSIYFKME NDLPQKIQCTLSNPLFNTTSSIIILTCIPSSGHSRHRVDDIEGRMDEPKSCDKTHTCPPCPAPPELLGGPSVFLFPPK PKDTLMISRTPPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKE YKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKT TPPVLDSDGSGFFLYSKLTVDKSRWQQGNVFCVMHEALHNHYTQKLSLSLSPGK

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.