

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7832: Recombinant Mouse Pulmonary Surfactant-associated Protein D/SP-D (C-6His)

Gene ID: 20390 **Uniprot ID:** P50404

Description

Source: Human Cells.

MW:36.7kD.

Recombinant Mouse Pulmonary Surfactant-associated Protein D is produced by our Mammalian expression system and the target gene encoding Ala20-Phe374 is expressed with a 6His tag at the C-terminus. SP-D (surfactant protein-D) is a 43 kDa member of the collectin family of innate immune modulators. Mouse SP-D cDNA encodes a 19 aa signal sequence and a 355 aa mature region with a 25 aa N-terminal linking-region, a 177 aa hydroxyproline and hydroxylysine collagen-like domain, a 46 aa coiled-coil segment, and a 106 aa, C-terminal collectin-like C-type lectin domain .SP-D is usually found as a glycosylated, disulfide-linked 150 kDa alpha -helical coiled-coil trimer with a "head" of three symmetrical CRDs . SP-D also binds SIRP alpha and the calreticulin/CD91 complex on macrophages.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: AEMKSLSQRSVPNTCTLVMCSPTENGLPGRDGRDGREGPRGEKGDPGLPGPMGLSGLQGPTGPVGPKGENG

SAGEPGPKGERGLSGPPGLPGIPGPAGKEGPSGKQGNIGPQGKPGPKGEAGPKGEVGAPGMQGSTGAKGSTG PKGERGAPGVQGAPGNAGAAGPAGPAGPQGAPGSRGPPGLKGDRGVPGDRGIKGESGLPDSAALRQQMEAL KGKLQRLEVAFSHYQKAALFPDGRSVGDKIFRTADSEKPFEDAQEMCKQAGGQLASPRSATENAAIQQLITAH NKAAFLSMTDVGTEGKFTYPTGEPLVYSNWAPGEPNNNGGAENCVEIFTNGOWNDKACGEORLVICEFVDHH

HHHH

Application Note

Storage condition:

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than $0.1 \text{ ng}/\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.