

32-7838: Recombinant Human Apolipoprotein C-II/APOC2 (C-6His)

Gene : APOC2

Gene ID : 344

Uniprot ID : P02655

Description

Source: E. coli.

MW :10kD.

Recombinant Human Apolipoprotein C-II is produced by our E.coli expression system and the target gene encoding Met1-Glu82 is expressed with a 6His tag at the C-terminus. APOC2 activates the lipoprotein lipase in capillaries, which hydrolyzes triglycerides and thus provides free fatty acids for cells. APOC2 is component of the very low density lipoprotein (VLDL) fraction in plasma. It is also an activator of several triacylglycerol lipases. The association of APOC2 with plasma chylomicrons, VLDL, and HDL is reversible, a function of the secretion and catabolism of triglyceride-rich lipoproteins, and changes rapidly. Defects in APOC2 are the cause of hyperlipoproteinemia type 1B (HLPP1B) which characterized by hypertriglyceridemia, xanthomas, and increased risk of pancreatitis and early atherosclerosis.

Product Info

Amount : 10 µg / 50 µg

Content : Supplied as a 0.2 µm filtered solution of 20mMPB,150mM NaCL,30%glycerol,pH7.2.

Storage condition : Store at -20°C, stable for 6 months after receipt. Please minimize freeze-thaw cycles.

Amino Acid : MTQPPQDDEMPSTLLTQVKESLSSYWESAKTAAQNLYEKTYLPAVDEKLRDLYSKSTAAMSTYTGIFTDQVLS
VLKGEELEHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.