

32-7839: Recombinant Human Dermatopontin/DPT/TRAMP (C-Fc-6His)(Discontinued)

Gene : DPT
Gene ID : 1805
Uniprot ID : Q07507

Description

Source: Human Cells.
MW :49.9kD.

Recombinant Human Dermatopontin is produced by our Mammalian expression system and the target gene encoding Gln19-Val201 is expressed with a Fc, 6His tag at the C-terminus. Dermatopontin, also known as Tyrosine-rich acidic matrix protein, TRAMP and DPT, is a secreted protein which belongs to the dermatopontin family. DPT is expressed in various tissues, such as fibroblasts, heart, skeletal muscle, brain and pancreas. It seems to mediate adhesion by cell surface integrin binding. DPT may serve as a communication link between the dermal fibroblast cell surface and its extracellular matrix environment. DPT can enhance TGFB1 activity through interaction with decorin. In addition, DPT accelerates collagen fibril formation, stabilizes collagen fibrils against low-temperature dissociation and inhibits cell proliferation.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : QYGDYGYYPYQQYHDYSDDGWVNLNRQGFYSYQCPQGQVIVAVRSIFSKEGSDRQWNYACMPTPQSLGEPTC
CWWEEINRAGMEWYQTCSNNGLVAGFQSRYSVLDREWQFYCCRYSKRCPYSCWLTIEYPGHYGEEMDMI
SYNYDYIRGATTTFSVERDRQWKFIMCRMTEYDCEFANVDDIEGRMDEPKSCDKTHTCPPCPAPELLGGP
SVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQ
DWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG
QPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGKHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.