

## 32-7844: Recombinant Mouse Granulocyte Colony-Stimulating Factor/G-CSF/CSF1(Discontinued)

**Gene :** Csf3  
**Gene ID :** 12985  
**Uniprot ID :** P09920

### Description

Source: Human Cells.

MW :18.8kD.

Recombinant Mouse Granulocyte Colony-Stimulating Factor is produced by our Mammalian expression system and the target gene encoding Val31-Ala208 is expressed. Granulocyte colony-stimulating factor (G-CSF) is a growth factor and an essential cytokine which belongs to the IL-6 superfamily. Granulocyte/macrophage colony-stimulating factors are cytokines that act in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and the monocytes-macrophages. G-CSF binding to its receptor G-CSF-R which belongs to the cytokine receptor type I family depends on the interaction of alpha-helical motifs of the former and two fibronectin type III as well as an immunoglobulin-like domain of the latter. G-CSF is a cytokine that have been demonstrated to improve cardiac function and perfusion in myocardial infarction. And it was initially evaluated as a stem cell mobilizer and erythropoietin as a cytoprotective agent.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** VPLVTVSALPPSLPLPRSFLLKSLEQVRKIQASGSVLLQLCATYKLCHPEELVLLGHSLGIPKASLSGCSSQALQQTQCLSQLHSGLCCLYQGLLQALSGISPALAPTLDLLQLDVANFATTIWQQMENLGVAPTQPTQSAMPAFTSAFQRRAGGVLAISYLQGFLETARLALHHLA

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.