

32-7845: Recombinant Human Interleukin-17B/IL-17B (C-Fc)

Gene : IL17B
Gene ID : 27190
Uniprot ID : Q9UHF5

Description

Source: Human Cells.
MW :45.1kD.

Recombinant Human Interleukin-17B is produced by our Mammalian expression system and the target gene encoding Gln21-Phe180 is expressed with a Fc tag at the C-terminus. Interleukin-17B (IL-17B) is a member of IL-17 cytokines family that plays important roles in host defence responses and inflammatory diseases. In addition to IL-17B, members of the IL-17 family include IL-17A, IL-17C, IL-17D, IL-17E and IL-17F. The six IL-17 cytokines are highly conserved at C terminus, and contain five spatially conserved cysteine residues that mediate dimerization. Expression of IL-17B protein has been reported in neurons, testis, ovary, stomach, pancreas, small intestine and chondrocytes. IL-17B binds to Interleukine-17 receptor B (IL-17 RB) and induces the production of inflammatory cytokines and the infiltration of inflammatory immune cells.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : QPRSPKSKRKGQGRPGPLAPGPHQVPLDLVSRMKPYARMEEYERNIEEMVAQLRNSSELAQRKCEVNLQLWM SNKRSLSPWGY SINHDPSRIPVDLPEARCLCLGCVNPFTMQEDRSMVSVFVSQVPVRRRLCPPPPRTGPCRQ RAVMETIAVGCTCIFIIEGRMDEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSH EDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKG QPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKS RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.