

32-7857: Recombinant Human Neuronal Cell Adhesion Molecule/NRCAM (C-Fc)(Discontinued)

Gene : NRCAM
Gene ID : 4897
Uniprot ID : Q92823

Description

Source: Human Cells.
MW :91.2kD.

Recombinant Human NRCAM is produced by our Mammalian expression system and the target gene encoding Gln25-Asn600 is expressed with a Fc tag at the C-terminus. Neuronal cell adhesion molecule(NRCAM) is a single-pass type I membrane protein ,containing 5 fibronectin type-III domains and 6 Ig-like C2-type (immunoglobulin-like) domains.It belongs to the immunoglobulin superfamily. NRCAM is engaged in such biological processes as axonal fasciculation, cell-cell adhesion, central nervous system development, clustering of voltage-gated sodium channels, neuron migration, positive regulation of neuron differentiation, regulation of axon extension, and synaptogenesis. It also may play a role in the molecular assembly of the nodes of Ranvier. NRCAM effects are also linked with different recognition processes and signal transduction pathways regulating cell differentiation, proliferation, or migration

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4 .
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : QMISALEVPLDPKLELDLVQPPTITQQSPKDYIIDPRENIVIQCEAKGKPPPSFSWTRNGTHFDIDKDPLVTMKPG TGTLIINIMSEGKAETYEGVYQCTARNERGA AVSNNIVVRPSRSP LWTKKLEPITLQSGQSLVLP CRPPIGLPPPII FWMDNSFQRLPQSERVSQGLNGDLYFSNVLPEDTREDYICYARFNHTQT IQKQPISVKVISAKSSRERPPPTFLT PEGNASNKEELRGNVLSLEICAEGLPTPIYWAKEDGMLPKNRTVYKNFEKTLQIIHVSEADSGNYQCI AKNALGA IHHTISVRVKAAPYWITAPQNLVLSPGEDGTLICRANGNPKPRISWLTNGVPIEAPDDPSRKIDGDTIIFSNVQER SSAVYQCNASNEYGYLLANAFVNVLAEP RILTPANTLYQVIANRPALLDCAFFGSP LPTIEWFKGAKGSALHEDI YVLHENGTL EIPVAQKDSTGTYTCVARNKLGMAKNEVHLEIKDATWIVKQPEYAVVQRGSMVSFECKVKHDHT LSLTVLWLKDNRELPSDERFTVDKDHLLVADVSDDDSGTYTCVANVDDIEGRMDEPKSCDKTHTCPPCPAPEL LGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTV LHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWE SNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNV FSCSVMHEALHNHYTQKSLSLSPGK

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.