

32-7865: Recombinant Mouse Collagen Alpha-1(XVIII) Chain/Endostatin (C-6His)

Gene : Col18a1
Gene ID : 12822
Uniprot ID : P39061

Description

Source: Human Cells.
MW :21.2kD.

Recombinant Mouse Collagen alpha-1(XVIII) chain is produced by our expression system and the target gene encoding His1591-Lys1774 is expressed Endostatin, an endogenous non-glycosylated inhibitor of endothelial cell proliferation and angiogenesis. It is produced and/or trimmed by metalloproteinases such as MMP-2 and MMP-9, and cathepsins S, B and L. The N-terminal ~27 aa of Endostatin appear to contain the majority of its activity. This region contains zinc binding sites that are thought to be critical for its anti-endothelial and anti-tumor effects, as well as multiple cleavage sites that, when used, can modify its activity. Mouse Endostatin shares 96% aa sequence identity with rat and 85-87% with human, bovine and equine Endostatin. It is predominantly expressed in liver, kidney, lung, skeletal muscle and testis. Endostatin inhibits endothelial cell growth by inducing cell cycle arrest in G1 phase and initiating apoptosis. It is also thought to down-regulate angiogenesis by blocking VEGF-induced endothelial cell migration. Endostatin may also be involved with down-regulation of angiogenesis after establishment of placental circulation in the pregnant uterus.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4 .
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : HTHQDFQPVLHLVALNTPLSGGMRGIRGADFQCFQQARAVGLSGTFRAFLSSRLQDLYSIVRRADRGSVPIVN LKDEVLSPSWDSLFSGSQGQLQPGARIFSFDRDVLRHPAWPQKSVVHGSDPSGRRLMESYCETWRTETTTG ATGQASSLLSGRLLLEQKAASCHNSYIVLCIENSFMTSFSKHHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.