

## 32-7872: Recombinant Mouse LAIR1 (C-6His)

**Gene :** Lair1  
**Gene ID :** 52855  
**Uniprot ID :** Q8BG84

### Description

Source: Human Cells.  
MW :14.4kD.

Recombinant Mouse Leukocyte-associated Immunoglobulin-like Receptor 1 is produced by our Mammalian expression system and the target gene encoding Gln22-Tyr141 is expressed with a 6His tag at the C-terminus. Leukocyte-associated Ig-like receptor-1 (LAIR-1) is an inhibitory receptor of the Ig superfamily that is structurally related to inhibitory members of KIR and ILT/CD85 families. It is expressed on immune cells, including NK cells, T cells, B cells, monocytes, immature neutrophils, dendritic cells and most thymocytes. The 253 amino acid (aa) type I transmembrane (TM) protein contains a 21 aa signal sequence, a 124 aa extracellular domain (ECD), a 20 aa TM domain and a 98 aa cytoplasmic domain. The ECD includes one C2-type Ig-like domain and two potential N-linked glycosylation sites. Tyrosine phosphorylation of two cytoplasmic ITIM motifs results in recruitment of phosphatases and down-regulation of signaling through activating receptors. LAIR1 shows high-affinity binding of collagens that results in inhibition of degranulation in a basophilic leukemia cell line.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** QEGSLPDITIFPNSSLMISQGTFTVVCSYSDKHDLYNMVRLEKDGSTFMEKSTEPYKTEDEFEIGPVNETITGHY  
SCIYSKGITWSERSKTLLELKVIKENVIQTPAPGPTSDTSWLKTYHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.