

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7885: Recombinant Human Interleukin-21/IL-21

Gene ID: 59067 Uniprot ID: Q9HBE4

Description

Source: E. coli. MW :15.6kD.

Recombinant Human Interleukin-21 is produced by our E.coli expression system and the target gene encoding Gln23-Ser155 is expressed. IL-21 induces the production of IgG1 and IgG3 in B-cells. IL-21 may promote the transition between innate and adaptive immunity. IL-21 may have a role in proliferation and maturation of natural killer cells in synergy with IL15. IL-21 stimulates interferon gamma production in T-cells and natural killer cells by synergy with IL15 and IL18. Dysregulation of IL-21 has a role in multiple immune-mediated diseases including lupus, psoriasis and chronic inflammatory diseases.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: MQGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWSAFSCFQKAQLKSANTGNNERIINVSIKK

LKRKPPSTNAGRRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.

Biological Activity: Measured by its ability to binding recombinant human IL2RG used funtional ELISA.