

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7892: Recombinant Human Ephrin-B1/EFNB1 (C-6His)

Gene ID: 1947
Uniprot ID: P98172

## **Description**

Source: Human Cells. MW :23.4kD.

Recombinant Human Ephrin-B1 is produced by our Mammalian expression system and the target gene encoding Leu28-Gly232 is expressed with a 6His tag at the C-terminus. Ephrin-B1, also named EFL-3, ELK ligand, EPH-related receptor tyrosine kinase ligand 2, is a single-pass type I membrane protein. It contains 1 ephrin RBD (ephrin receptor-binding) domain and belongs to the ephrin family. Ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. All ephrins share a conserved extracellular sequence, which most likely corresponds to the receptor-binding domain. Ephrin-B1 has been shown to bind EphA3, EphB1, EphB2, EphB3, and EphB4. The extracellular domains of human and mouse ephrin-B1 share 94% amino acid identity.

## **Product Info**

**Amount:**  $10 \mu g / 50 \mu g$ 

**Content:** Lyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid: LAKNLEPVSWSSLNPKFLSGKGLVIYPKIGDKLDIICPRAEAGRPYEYYKLYLVRPEQAAACSTVLDPNVLVTCNR

PEQEIRFTIKFQEFSPNYMGLEFKKHHDYYITSTSNGSLEGLENREGGVCRTRTMKIIMKVGQDPNAVTPEQLTT

SRPSKEADNTVKMATQAPGSRGSLGDSDGKHETVNQEEKSGPGASGGSSGDPDGVDHHHHHH

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin**: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.