

32-7900: Recombinant Human Nucleotide Exchange Factor SIL1/SIL1 (C-6His)(Discontinued)

Gene : SIL1
Gene ID : 64374
Uniprot ID : Q9H173

Description

Source: Human Cells.
MW :49.8kD.

Recombinant Human Nucleotide exchange factor SIL1 is produced by our Mammalian expression system and the target gene encoding His32-Arg461 is expressed with a 6His tag at the C-terminus. Nucleotide exchange factor SIL1, also named BiP-associated protein, is a member of the SIL1 family. It is a resident endoplasmic reticulum (ER), N-linked glycoprotein with an N-terminal ER targeting sequence, 2 putative N-glycosylation sites, and a C-terminal ER retention signal. It is highly expressed in tissues which produce large amounts of secreted proteins such as kidney, liver and placenta. This protein functions as a nucleotide exchange factor for another unfolded protein response protein. Mutations in this gene have been associated with Marinesco-Sjogren syndrome.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : HQNLKEFALTNPEKSSTKETERKETAEEELDAEVLEVFHPTHEWQALQPGQAVPAGSHVRLNLQTGEREAKL QYEDKFRNNLKGKRLDINTNTYTSQDLKSALAKFKEGAEMESSKEDKARQAEVKRFRPIEELKKDFDELNVVIE TDMQIMVRLINKFNSSSSSLEEKIAALFDLEYVHQM DNAQDLLSFGGLQVVINGLNSTEPLVKEYAAFVLGAAF SSNPKVQVEAIEGGALQKLLVILATEQPLTAKKKVLFALCSLLRHFPYAQRQFLKLGGLQVLR TLVQEKGT EVLAV RVVTL LYDLVTEKMF AEEEAELTQEMSPEKLQYRQVHLLPGLWEQGWCEITAHLLALPEHDAREKVLQTLGV LLTTCRDRYRQDPQLGR TLASLQAEYQVLASLELQDGEDEGYFQELLGSVNSLLKELRVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.