

## 32-7903: Recombinant Human Pregnancy-Specific beta-1-Glycoprotein 1/PSG1 (C-6His)

**Gene :** PSG1  
**Gene ID :** 5669  
**Uniprot ID :** P11464

### Description

Source: Human Cells.  
MW :44.5kD.

Recombinant Human Pregnancy-specific beta-1-glycoprotein 1 is produced by our Mammalian expression system and the target gene encoding Gln35-Pro419 is expressed with a 6His tag at the C-terminus. Pregnancy-specific beta-1-glycoprotein 1(PSBG-1 for short), also named CD66 antigen-like family member F, Fetal liver non-specific cross-reactive antigen 1/2, PSG95, Pregnancy-specific beta-1 glycoprotein C/D, is a secreted protein which belongs to the immunoglobulin superfamily, CEA family. It contains 3 Ig-like C2-type (immunoglobulin-like) domains and 1 Ig-like V-type (immunoglobulin-like) domain. The human placenta is a multihormonal endocrine organ that produces hormones, enzymes, and other molecules that support fetal survival and development.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** QVTIEAEPTKVSEKDVLLLVHNLQNLTYIWKYQMRDLYHYITSYVVDGEIIYGPAYSGRETAYSNASLLIQ NVTREDAGSYTLHIKGGDGTGRVTRFTFLHLETPKPSISSNLNPRETMEAVSLTCDPETPDASYLWWMNG QSLPMTSHSLKLSLSETNRTLFLGLVTKYTAGPYECEIRNPVSASRSDPVTNLNLLPKLPKYITINNLNPRENKDVLNFT CEPKSENYTYIWWLNGQSLPVSPRVKRPNIENRILIPSVTRNETGPYQCEIRDYGGIRSDPVTNLNVLYGPDLPRIY PSFTYYRSGEVLYLSCSADSNPQAQYSWTINEKFQLPGQKLFIRHITTKHSGLYVCSVRNSATGKESKSMTEVEV SDWTVPVDHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.