

## 32-7929: Recombinant Human Follistatin-Related Protein 1/FSTL1 (C-6His)(Discontinued)

**Gene :** FSTL1  
**Gene ID :** 11167  
**Uniprot ID :** Q12841

### Description

Source: Human Cells.  
MW :33.5kD.

Recombinant Human Follistatin-like Protein 1 is produced by our Mammalian expression system and the target gene encoding Glu21-Ile308 is expressed with a 6His tag at the C-terminus. Follistatin-Related Protein 1 (FSTL1) is a secreted protein that contains two EF-hand domains, one follistatin-like domain, one Kazal-like domain, and one VWFC domain. Its functional significance in physiological and pathological processes is not completely understood. However, FSTL1 is thought to modulate the action of some growth factors on cell proliferation and differentiation. FSTL1 maybe an autoantigen associated with rheumatoid arthritis.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** EEELRSKSKICANVFCGAGRECAVTEKGEPTCLCIEQCKPHKRPVCGSNGKTYLNHCELHRDACTLGSKIQVDY DGHCKEKKSVSPSASPVCYQSNRDELRRRIQWLEAEIIPDGWFSKGSNYSEILDYFKNFDNGDSRLDSSEFL KFVEQNETAINITYPDQENKLLRGLCVDALIELSDENADWKLSFQFLKCLNPSFNPPEKKCALEDETYADGA ETEVDCNRCVCACGNWVCTAMTCDGKNQKGAQTQTEEMTRYVQELQKHQETAETKRVSTKEIHSHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.