

32-7938: Recombinant Human Interleukin-10/IL-10(Discontinued)

Gene : IL10
Gene ID : 3586
Uniprot ID : P22301

Description

Source: Human Cells.
MW :19.6kD.

Recombinant Human Interleukin-10 is produced by our Mammalian expression system and the target gene encoding Ser19-Asn178 is expressed. Interleukin 10(IL10), also known as cytokine synthesis inhibitory factor (CSIF), is a secreted protein and belongs to the IL-10 family. IL-10 is secreted by many activated hematopoietic cell types as well as hepatic stellate cells, keratinocytes, and placental cytotrophoblasts. IL-10 is an anti-inflammatory TH2 cytokine that has a critical role in limiting the immune response to pathogens to prevent host damage. As IL-10 is produced in several T helper populations, it is proposed that it provides a feedback loop to limit the effector functions of macrophages and DCs on T cells. Once expressed, IL-10 signals through the IL-10 receptor (IL-10R) to activate STAT3. As IL-10 is a strong inhibitor of inflammation, it has become a viable biomarker for various diseases and conditions as well as a therapeutic molecule for certain conditions. In addition to elevated levels in parasitic infection, high expression levels of IL-10 are also found in retroviral infections inducing immunodeficiency. The immunosuppressive properties of IL-10 suggest a possible clinical use of IL-10 in suppressing rejections of grafts after organ transplantations

Product Info

Amount : 10 / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : SPGQGTQSENSCTHFPGNLPLMLRDLRDAFSRVKTFQMKDQLDNLKESLLEDFKGYLGCQALSEMIQFYL
EEVMPQAENQDPDIKAHVNSLGENLKLRLRLRCHRFLPCENKSKAVEQVKNAFNKLQEKGIYKAMSEFDIFIN
YIEAYMTMKIRN

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.