

## 32-7941: Recombinant Human Glial Cell Line-Derived Neurotrophic Factor/GDNF (C-Fc-6His)(Discontinued)

**Gene :** GDNF  
**Gene ID :** 2668  
**Uniprot ID :** P39905

### Description

Source: Human Cells.

MW :49.6kD.

Recombinant Human GDNF is produced by our Mammalian expression system and the target gene encoding Phe20-Ile211 is expressed with a Fc, 6His tag at the C-terminus. Glial Cell Line-Derived Neurotrophic Factor (GDNF) is a disulfide-linked homodimeric glycoprotein that belongs to the TGF- beta superfamily. It has been shown to promote the survival of various neuronal subpopulations in both the central as well as the peripheral nervous systems at different stages of their development. Human GDNF cDNA encodes a 211 amino acid residue prepropeptide that is processed to yield a dimeric protein. Mature human GDNF was predicted to contain two 134 amino acid residue subunits. Cells known to express GDNF include Sertoli cells, type 1 astrocytes, Schwann cells, neurons, pinealocytes and skeletal muscle cells. Mutations in this gene may be associated with Hirschsprung disease.

### Product Info

**Amount :** 6His) / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** FPLPAGKRPPEAPAEDRSLGRRRAPFALSSDSNMPEDYPDQFDDVMDFIQATIKRLKRSPDKQMAVLPRRERN  
RQAAAANPENSRRGKRRRQGRGKNRGCVLTAIHLNVTDLGLGYETKEELIFRYCSGSCDAAETTYDKILKNLSRN  
RRLVSDKVGQACCRPIAFDDDLFLDDNLVYHILRKHSKRRCGIVDDIEGRMDEPKSCDKTHTCPPCPAPPELL  
GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVL  
HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWES  
NGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQGNVFSVMSVHEALHNHYTQKSLSLSPGKHHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.