

32-7949: Recombinant Human Leukocyte Mono Ig-Like Receptor 2/LMIR2/CD300C (C-Fc)(Discontinued)

Gene : CD300C
Gene ID : 10871
Uniprot ID : Q08708

Description

Source: Human Cells.

MW :44.9kD.

Recombinant Human LMIR2 is produced by our Mammalian expression system and the target gene encoding Gly21-Arg183 is expressed with a Fc tag at the C-terminus. CD300C is a single-pass type I membrane protein which belongs to the immunoregulatory signaling (IRS) family. CD300C contains one Ig-like V-type domain and is present on the surface of natural killer cells, granulocytes, most myeloid cells, dendritic cells, and a subpopulation of T and B lymphocytes. The CD300C (CMRF-35A) and CD300A (CMRF-35H) molecules are homologous leukocyte surface proteins. CD300a and CD300C play an important role in the cross-regulation of TNF-alpha and IFN-alpha secretion from pDCs. CD300A and CD300C are indistinguishable on the surface of NK cells. The ligand for CD300C is presently unknown.

Product Info

Amount : Fc) / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : GYFPLSHPMTVAGPVGGSLSVQCRYEKEHRTLNFVCRPPQILRCDKIVETKGSAGKRNGRVSIRDSPANLSFT VTLENLTEEDAGTYWCGVDPWLRDFHDPIVEVEVSVPAGTTTASSPQSSMGTS GPPTKLPVHTWPSVTRKD SPEPSPHPGSLFSNVRVDDIEGRMDEPKSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVD VSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISK AKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSSFFLYSKLTV DKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.