

## 32-7957: Recombinant Mouse Cathepsin L/CTSL (C-6His)(Discontinued)

**Gene :** Ctsl  
**Gene ID :** 13039  
**Uniprot ID :** P06797

### Description

Source: Human Cells.  
MW :36.8kD.

Recombinant Mouse Cathepsin L is produced by our Mammalian expression system and the target gene encoding Thr18-Asn334 is expressed with a 6His tag at the C-terminus. Mouse Cathepsin L is a lysosomal cysteine proteinase which is a member of the peptidase C1 family. This protein is a dimer composed of disulfide-linked heavy and light chains, both produced from a single protein precursor. Cathepsin L plays a major role in intracellular protein catabolism. Its substrates include collagen and elastin, as well as alpha-1 protease inhibitor, a major controlling element of neutrophil elastase activity. Cathepsin L has been implicated in several pathologic processes, including myofibril necrosis in myopathies and in myocardial ischemia, and in the renal tubular response to proteinuria.

### Product Info

**Amount :** 6His) / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** TPKFDQTFSAEWHQWKSTHRRLYGTNEEEWRRRAIWEKNMRMIQLHNGEYSNGQHGFSMEMNAFGDMTNEE  
FRQVVNGYRHQKHKKGRLFQEPLMLKIPKSVDWREKGCVTPVKNQGCQSCWAFSASGCLEGQMFLKTGKLI  
SLSEQNLVDCSHAQGNQGCNGLMDFAFQYIKENGGLDSEESYPYEAKDGSCYRAEFVANDTGFDIPQQ  
EKALMKAVATVGPISVAMDASHPSLQFYSSGIYEPNCSSKNLDHGVLLVGYEGTDSNKNKYWLKNSWGS  
EWGMEGYIKAKDRDNHCGLATAASYPVVNVDDHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.