

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7960: Recombinant Mouse Cathepsin E/CTSE (C-6His)(Discontinued)

**Gene ID:** 13034 **Uniprot ID:** P70269

## **Description**

Source: Human Cells.

MW:37kD.

Recombinant Mouse Cathepsin E is produced by our Mammalian expression system and the target gene encoding Ser60-Pro397 is expressed with a 6His tag at the C-terminus. Cathepsin E is encoded by the ctse gene, exists in the homodimer forms, belongs to the peptidase A1 family. Cathepsin E high expressed in the stomach, clara cells and alveolar macrophages of lung, brain microglia, spleen and activated B-lymphocytes. Cathepsin E may involve in the processing of antigenic peptides during MHC class II-mediated antigen presentation, play a role in activation-induced lymphocyte depletion in the thymus, and in neuronal degeneration and glial cell activation in the brain.

## **Product Info**

**Amount:** 6His) / 50 μg

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: SCNVYSSVNEPLINYLDMEYFGTISIGTPPQNFTVIFDTGSSNLWVPSVYCTSPACKAHPVFHPSQSDTYTEVGN

HFSIQYGTGSLTGIIGADQVSVEGLTVDGQQFGESVKEPGQTFVNAEFDGILGLGYPSLAAGGVTPVFDNMMA QNLVALPMFSVYLSSDPQGGSGSELTFGGYDPSHFSGSLNWIPVTKQAYWQIALDGIQVGDTVMFCSEGCQAI VDTGTSLITGPPDKIKQLQEAIGATPIDGEYAVDCATLDTMPNVTFLINEVSYTLNPTDYILPDLVEGMQFCGSGF

QGLDIPPPAGPLWILGDVFIRQFYSVFDRGNNQVGLAPAVPVDHHHHHH

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin**: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.