

## 32-7965: Recombinant Human Clusterin/ApoJ (C-Fc-6His)(Discontinued)

**Gene :** CLU  
**Gene ID :** 1191  
**Uniprot ID :** P10909

### Description

Source: Human Cells.  
MW :78kD.

Recombinant Human Clusterin is produced by our Mammalian expression system and the target gene encoding Asp23-Glu449 is expressed with a Fc, 6His tag at the C-terminus. Clusterin is a secreted protein which belongs to the Clusterin family. Clusterin is expressed in adult testis, heart, ovary, adrenal gland, brain and liver. Clusterin has been suggested to be involved in several basic biological events such as cell death, tumor progression, and neurodegenerative disorders. In addition, Clusterin is up/ down regulated on the mRNA or protein level in many pathological and clinically relevant situations including cancer, organ regeneration, infection, Alzheimer disease, retinitis pigmentosa, myocardial infarction, renal tubular damage, autoimmunity and others.

### Product Info

**Amount :** 6His) / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** DQTVSDNELQEMSNQGSKYVNKEIQNAVNGVKQIKTLIEKTNEERKTLLSNLEEAKKKKEDALNETRESETKLKE  
LPGVCNETMMALWEECKPCLKQTCMKFYARVCRSGSLVGRQLEEFNLQSSPFYFWMNGDRIDSLLENDRQ  
QTHMLDVMQDHFSSRASSIIDELFQDRFFTREPDQTYHYLPFSLPHRRPHFFPKSRIVRSLMPFSPYEPLNFHAM  
FQPFLEMIHEAQQAMDIHFHSPAQHPPTFEIREGDDDRVTVCREIRHNSTGCLRMKDQCDKCREILSVDCTNN  
PSQAKLRRELDSELQVAERLTRKYNELLKSYQWKMLNTSSLEQLNEQFNWVSRLANLTQGEDQYYLRVTTVA  
SHTSDSDVPSGVTEVVVKLFDSDPITVTPVEVSRKNPKFMETVAEKALQEYRKKHREEVDDIEGRMDEPKSCD  
KTHTCPPELPGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQ  
YNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLV  
KGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSL  
SLSPGKHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.