

## 32-7974: Recombinant Human NKG2D Ligand 2/NKG2DL2/N2DL2 (C-Fc)(Discontinued)

**Gene :** ULBP2  
**Gene ID :** 80328  
**Uniprot ID :** Q9BZM5

### Description

Source: Human Cells.  
MW :51.4kD.

Recombinant Human NKG2D ligand 2 is produced by our Mammalian expression system and the target gene encoding Gly26-Ser217 is expressed with a Fc tag at the C-terminus. NKG2D Ligand 2 (N2DL2) is a member of a family of cell-surface proteins. N2DL2 function as ligands for human cytomegalovirus glycoprotein UL16. N2DL2 is anchored to the membrane via a GPI-linkage. N2DL2 is bind to human NKG2D, an activating receptor expressed on NK cells, NKT cells, T cells. Engagement of NKG2D results in the activation of cytolytic activity and cytokine production by these effects cells. The ULBPs are expressed on some tumor cells and have been implicated in tumor surveillance.

### Product Info

**Amount :** Fc) / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** GRADPHSLCYDITVIPKFRPGPRWCAVQGQVDEKTFLLHYDCGNKTVTPVSPGLGKKNVTTAWKAQNPVLEVV  
DILTEQLRDIQLENYTPKEPLTLQARMSCEQKAEGHSSGSWQFSFDGQIFLLFDSEKRMWTTVHPGARKMKEK  
WENDKVVAMSFHYFSMGDCIGWLEDFLMGMDSTLEPSAGAPLAMSSVDDIEGRMDEPKSCDKTHTCPPCPA  
PELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSV  
LTVLHQDWLNGKEYKCKVSNKALPAPIEKTKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVE  
WESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVDFCSVMHEALHNHYTQKSLSLSPGK

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.