

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7987: Recombinant Mouse Interleukin-13/IL-13 (Ser26-Phe131,C-6His)(Discontinued)

Gene ID: 16163 **Uniprot ID:** P20109

Description

Source: Human Cells.

MW:12.7kD.

Recombinant Mouse Interleukin-13 is produced by our Mammalian expression system and the target gene encoding Ser26-Phe131 is expressed with a 6His tag at the C-terminus. Mouse interleukin 13 (mIL-13) is a pleiotropic cytokine produced by activated Th2 cells. IL-13 induces B cell proliferation and immunoglobin production. It contains a four helical bundle with two internal disulfide bonds. Mouse IL13 shares 58% sequence identity with human protein and exhibits cross-species activity. IL13 signals via receptor IL13R (type2, IL4R) and activates STAT-6. IL13 initially binds IL-13Ra1 with low affinity and triggers association of IL4Ra, generating a high affinity heterodimeric receptor IL13R and eliciting downstream signals. IL13 also binds IL-13Ra2 with high affinity, which plays a role in a negative feedback system of IL13 signaling. IL13 is an important mediator of allergic inflammation and disease.

Product Info

Amount: 6His) / 50 μg

Content: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

Storage condition : Reconstituted protein solution can be stor samples are stable at -20°C for 3 months.

Amino Acid: SVSLPLTLKELIEELSNITQDQTPLCNGSMVWSVDLAAGGFCVALDSLTNISNCNAIYRTQRILHGLCNRKAPTTV

SSLPDTKIEVAHFITKLLSYTKQLFRHGPFVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.