

32-7988: Recombinant Human Activin Receptor 2B/Activin RIIB/ACVR2B (C-Fc-6His)(Discontinued)

Gene : ACVR2B
Gene ID : 93
Uniprot ID : Q13705

Description

Source: Human Cells.

MW :41.3kD.

Recombinant Human Activin Receptor IIB is produced by our Mammalian expression system and the target gene encoding Ser19-Thr134 is expressed with a Fc, 6His tag at the C-terminus. Activin proteins that belong to the transforming growth factor-beta (TGF- beta) superfamily, exert their biological actions by binding to heteromeric receptor complexes of type I and type II serine/threonine kinase receptors. On ligand binding, type I and II receptors form a stable complex, resulting in phosphorylation of type I receptors by type II receptors with constitutive kinase activity, and subsequently initiates the activation of downstream molecules including the endogenous Smads. ActRIIB, also known as ActRIIB, is a type II receptor containing an extracellular domain (ECD), a transmembrane segment, and a cytoplasmic region that includes the kinase domain. ActRIIB is a receptor for activin A, activin B and inhibin A. Multiple ActRIIB isoforms can also be generated, which bind activin isoforms with different affinities.

Product Info

Amount : 6His) / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : SGRGEAETRECIYYNANWELERTNQSLERCEGEQDKRLHCYASWRNSSGTIELVKKGCWLDDFNCDYDRQEC VATEENPQVYFCCCEGNFCNERFTHLPEAGGPEVTYEPPPTAPTVDIEGRMDPEKSCDKTHTCPPCPAPELLG GPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLH QDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESN GPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQGNVFSCSVMHEALHNHYTQKSLSLSPGKHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.