

32-7990: Recombinant Human CD27/TNFRSF7 (C-Fc-6His)(Discontinued)

Gene : CD27
Gene ID : 939
Uniprot ID : P26842

Description

Source: Human Cells.
MW :47.2kD.

Recombinant Human CD27 is produced by our Mammalian expression system and the target gene encoding Thr21-Ile192 is expressed with a Fc, 6His tag at the C-terminus. CD27 antigen is also known as CD27L receptor, T-cell activation antigen CD27, Tumor necrosis factor receptor superfamily member 7, T14 and TNFRSF7. In humans, it is encoded by the CD27 gene. CD27 is a single-pass type I membrane protein with 3 TNFR-Cys repeats. It is a member of the TNF-receptor superfamily and is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. It plays a role in survival of activated T-cells and apoptosis through association with SIVA1.

Product Info

Amount : 6His) / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM Tris,150mM NaCl, pH 8.0.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : TPAPKSCPERHYWAQGLCCQMCEPGTFLVKDCDQHRKAAQCDCIPGVSFSPDHHTRPHCESCRHCNSGLLVRNCTITANAECACRNGWQCRDKECTCDPLPNPSLTARSSQALSPHPQPTHLPYVSEMLEARAGHMQTLADFRQLPARTLSTHWPPQRSRLCSSDFIRIVDDIEGRMDEPKSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN KALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDS DGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGKHHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.