

## 32-8001: Recombinant Human T Cell Immunoglobulin and Mucin Domain-3/TIM-3/HAVCR2 (C-Fc-6His)(Discontinued)

**Gene :** HAVCR2

**Gene ID :** 84868

**Uniprot ID :** Q8TDQ0

### Description

Source: Human Cells.

MW :47.7kD.

Recombinant Human TIM-3 is produced by our Mammalian expression system and the target gene encoding Ser22-Arg200 is expressed with a Fc, 6His tag at the C-terminus. Hepatitis A virus cellular receptor 2(HAVCR2) is a single-pass type I membrane protein and it contains 1 Ig-like V-type (immunoglobulin-like) domain. The protein belongs to the immunoglobulin superfamily, and TIM family of proteins. The protein regulates macrophage activation. It inhibits T-helper type 1 lymphocyte (Th1)-mediated auto- and alloimmune responses and promotes immunological tolerance. It may be also involved in T-cell homing and it is receptor for LGALS9. CD4 (MIM 186940)-positive T helper lymphocytes can be divided into types 1 (Th1) and 2 (Th2) on the basis of their cytokine secretion patterns. Th1 cells and their associated cytokines are involved in cell-mediated immunity to intracellular pathogens and delayed-type hypersensitivity reactions, whereas Th2 cells are involved in the control of extracellular helminthic infections and the promotion of atopic and allergic diseases. The 2 types of cells also cross-regulate the functions of the other. TIM3 is a Th1-specific cell surface protein that regulates macrophage activation and enhances the severity of experimental autoimmune encephalomyelitis in mice.

### Product Info

**Amount :** 6His) / 50 µg

**Content :** Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

**Amino Acid :** SEVEYRAEVEGQNAYLPCFYTPAAPGNLVPVCWGKGACPVFECGNVVLRTDERDVNYWTSRYWLNDFRKGDVSLTIENVTLADSGIYCCRIQIPGIMNDEKFNKLVIKPAKVTPAPTLQRDFTAAPRMLTTRGHGPAETQTLGSLPDINLTQISTLANELRDSRLANDLRDSGATIRVDDIEGRMDEPKSCDKHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSSFFLYSKLTVDKSRWQQGNVVFSCSVMEALHNHYTQKSLSLSPGKHHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.