

32-8002: Recombinant Human Interleukin-18/IL-18/IL-1F4 (C-6His)(Discontinued)

Gene : IL18
Gene ID : 3606
Uniprot ID : Q14116

Description

Source: Human Cells.
MW :23.3kD.

Recombinant Human Interleukin-18 is produced by our Mammalian expression system and the target gene encoding Tyr37-Asp193 is expressed with a 6His tag at the C-terminus. Interleukin-18 is a secreted protein and it belongs to the IL-1 family. IL-18 is a proinflammatory cytokine and produced by macrophages and other cells. This cytokine can induce the IFN-gamma production of T cells. The combination of this cytokine and IL12 has been shown to inhibit IL-4 dependent IgE and IgG1 production, and enhance IgG2a production of B cells. IL-18 binding protein (IL18BP) can specifically interact with this cytokine, and thus negatively regulate its biological activity. After stimulation with IL-18, natural killer (NK) cells and certain T cells release another important cytokine called interferon-gamma (IFN-gamma) or type II interferon that plays an important role in activating the macrophages or other cells.

Product Info

Amount : 6His) / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.
Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.
Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : YFGKLESKLSVIRNLNDQVLFIDQGNRPLFEDMTDSDCRDNAPRTIFIISMKYKDSQPRGMAVTISVKCEKISTLSC
ENKIISFKEMNPPDNIKDTKSDIIFQRSVPGHDNKMQFESSYEGYFLACEKERDLFKLILKKEDELGDRSIMFTV
QNEDVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.