

32-8004: Recombinant Mouse Ephrin-A1/EFNA1 (C-Fc-6His)(Discontinued)

Gene : Efn1
Gene ID : 13636
Uniprot ID : P52793

Description

Source: Human Cells.
MW :47.3kD.

Recombinant Mouse Ephrin-A1 is produced by our Mammalian expression system and the target gene encoding Asp19-Ser182 is expressed with a Fc, 6His tag at the C-terminus. Ephrin-A1 is a cell membrane protein and contains 1 ephrin RBD (ephrin receptor-binding) domain. EFNA1 belongs to the ephrin (EPH) family. The ephrins and EPH-related receptors comprise the largest subfamily of receptor protein-tyrosine kinases and have been implicated in mediating developmental events, especially in the nervous system and in erythropoiesis. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. This gene encodes an EFNA class ephrin which binds to the EPHA2, EPHA4, EPHA5, EPHA6, and EPHA7 receptors. Two transcript variants that encode different isoforms were identified through sequence analysis. It belongs to the ephrin family and contains 1 ephrin RBD (ephrin receptor-binding) domain.

Product Info

Amount : 6His) / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : DRHIVFWNSSNPKFREEDYTVHVQLNDYLDIICPHYEDDSVADAAMERYTYLYMVEHQEYVACQPQSKDQVRW
NCNRPSAKHGPEKLSEKFQRFTPFILGKEFKEGHSYYYISKPIYHQESQCLKLVTVNGKITHNPQAHVNPQEKRL
QADDPEVQVLHSGYSVDDIEGRMDEPKSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVV
DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTIS
KAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLT
VDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGKHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.