

32-8005: Recombinant Mouse TWEAK Receptor/TWEAK R/TNFRSF12A (C-Fc)(Discontinued)

Gene : Tnfrsf12a

Gene ID : 27279

Uniprot ID : Q9CR75

Description

Source: Human Cells.

MW :32.6kD.

Recombinant Mouse TWEAK Receptor is produced by our Mammalian expression system and the target gene encoding Glu28-Trp79 is expressed with a Fc tag at the C-terminus. Tumor necrosis factor receptor superfamily member 12A(Tnfrsf12a) is a single-pass type I membrane protein and contains 1 TNFR-Cys repeat. It is weak inducer of apoptosis in some cell types. It promotes angiogenesis and it is the proliferation of endothelial cells. It may modulate cellular adhesion to matrix proteins. TNFR binds specifically to tumor necrosis factor (TNF) and blocks its interaction with cell surface TNF receptors. TNF is a naturally occurring cytokine that is involved in normal inflammatory and immune responses. It plays an important role in the inflammatory processes of rheumatoid arthritis (RA), polyarticular-course juvenile rheumatoid arthritis (JRA), and ankylosing spondylitis and the resulting joint pathology. In addition, TNF plays a role in the inflammatory process of plaque psoriasis. Elevated levels of TNF are found in involved tissues and fluids of patients with RA, psoriatic arthritis, ankylosing spondylitis (AS), and plaque psoriasis. Two distinct receptors for TNF (TNFRs), a 55 kilodalton protein (p55) and a 75 kilodalton protein (p75), exist naturally as monomeric molecules on cell surfaces and in soluble forms. Biological activity of TNF is dependent upon binding to either cell surface TNFR.

Product Info

Amount : Fc) / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : EQAPGTSPCSSGSSWSADLDKCMDCASCARP HSDFC LGCAAAPAHFRLLWVDDIEGRMDEPKSCDKTHT
CPPCPAPPELLGGPSVFLFPPKPKD TLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST
YRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYP
SDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNV FSCVMHEALHNHYTQKLSLSLSPG
K

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.