

32-8009: Recombinant Mouse Death Receptor 6/DR6/TNFRSF21/CD358 (C-Fc-6His)(Discontinued)

Gene : Tnfrsf21
Gene ID : 94185
Uniprot ID : Q9EPU5

Description

Source: Human Cells.

MW :64.7kD.

Recombinant Mouse Death Receptor 6 is produced by our Mammalian expression system and the target gene encoding Gln42-His349 is expressed with a Fc, 6His tag at the C-terminus. Tumor necrosis factor receptor superfamily member 21(DR6) is a single-pass type I membrane protein and contains 1 death domain and 4 TNFR-Cys repeats. The protein may activate NF-kappa-B and promote apoptosis and it may activate JNK and be involved in T-cell differentiation. It is required for both normal cell body death and axonal pruning. Trophic-factor deprivation triggers the cleavage of surface APP by beta-secretase to release sAPP-beta which is further cleaved to release an N-terminal fragment of APP (N-APP). N-APP binds TNFRSF21 triggering caspase activation and degeneration of both neuronal cell bodies (via caspase-3) and axons (via caspase-6).

Product Info

Amount : 6His) / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : QPEQKTLSPGTYRHVDRITGQVLTCDKCPAGTYVSEHCTNMSLRVCSGPCAGTFTRHENGIERCHDCSQPCP
WPMIERLPCAALTDREICPPGMYQSNGTCAPHTVCPVGGVGRKKGTENEDVRCKQCARGTFSDVPSSVMKC
KAHTDCLGQNLVVKPGTKETDNVCGMRLFFSSTNPPSSGTVTFSHPEHMESHDPSSSTYEPQGMNSTDSNS
TASVRTKVPSEIEEGTVPDNTSSTSGKEGTNRTPNPPQVTHQQAPHHRHILKLLPSSMEATGEKSSTAIAKAPKR
GHPRQNAHKHFDINEHVDDIEGRMDEPKSCDKHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVV
DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTIS
KAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLT
VDKSRWQQGNVFSCSVMHEALHNHYTQKLSLSLSPGKHHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.