

32-8011: Recombinant Human CD28/TP44 (C-Fc)(Discontinued)

Gene : CD28
Gene ID : 940
Uniprot ID : P10747

Description

Source: Human Cells.
MW :42.3kD.

Recombinant Human CD28 is produced by our Mammalian expression system and the target gene encoding Asn19-Pro152 is expressed with a Fc tag at the C-terminus. T-cell-specific surface glycoprotein CD28(CD28) is a single-pass type I membrane protein which contains one Ig-like V-type (immunoglobulin-like) domain. It belongs to the immunoglobulin(Ig) superfamily. CD28 is one of the molecules expressed on T cells that provide co-stimulatory signals, which are required for T cell activation. CD28 co-stimulation is necessary for CD4 positive T-cell proliferation and survival, interleukin-2 production, and T-helper type-2 development. Human post-thymic regulatory T cells require CD28 co-stimulation to expand and maintain potent suppressive function in vivo. Apoptosis plays a key role in the age-related decline of CD28 expression and in immunosenescence. CD28 is the receptor for CD80 (B7.1) and CD86 (B7.2). When activated by Toll-like receptor ligands, the CD80 expression is upregulated in antigen presenting cells (APCs). The CD86 expression on antigen presenting cells is constitutive. CD28 is the only B7 receptor constitutively expressed on naive T cells.

Product Info

Amount : Fc) / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : NKILVKQSPMLVAYDNAVNLSCYKSYNLF SREFRASLHKGLDSAVEVCVVYGNYSQQLQVYSKTGFNCDGKLG NESVTFYLQNLVYNQTDIYFCKIEVYPPPYLDNEKSNGTIIHVKGKHLCPSPFLPGPSKPVDDIEGRMDEPKSCD KTHTCPPEPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQ YNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLV KGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNV FSCSVMHEALHNHYTQKSL SLSPGK

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.