

## 32-8020: Recombinant Human Receptor Tyrosine-Protein Kinase ErbB-3/HER3 (C-Fc)(Discontinued)

**Gene :** ERBB3  
**Gene ID :** 2065  
**Uniprot ID :** P21860

### Description

Source: Human Cells.

MW :61.6kD.

Recombinant Human ErbB3 is produced by our Mammalian expression system and the target gene encoding Ser20-Cys331 is expressed with a Fc tag at the C-terminus. Receptor tyrosine-protein kinase erbB-3 is an enzyme that in humans is encoded by the ERBB3 gene. This gene encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases. ERBB3 belongs to the protein kinase superfamily, tyr protein kinase family and EGF receptor subfamily. It contains 1 protein kinase domain and it is expressed in Epithelial tissues and brain. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However, it does form heterodimers with other EGF receptor family members which do have kinase activity.

### Product Info

**Amount :** Fc) / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** SEVGNSQAVCPGTLNGLSVTGDAENQYQTLTKLYERCEVVMGNLEIVLTGHNADLSFLQWIREVTGYVLVAMN EFSTLPLPNLRVVRGTQVYDYGKFAIFVMLNYNTNSSHALRQLRLTQLTEILSGGVYIEKNDKLCHMDTIDWRDIV RDRDAEIVVKDNGRSCPPCHEVCKGRCWGPGEEDCQTLTKTICAPQCNGHCFGNPNQCCHDECAGGCSGP QDTDCFACRHFNDSGACVPRCPQPLVYNKLTQLEPNPHTKYQYGGVVCVASCPhNFVVDQTSVCRACPPDKM EVDKNGLKMCEPCGGLCPKAFVDDIEGRMDEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAP IEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFLL YSKLTVDKSRWQQGNVVFCSVMHEALHNHYTQKSLSLSPGK

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.